arXiv:1610.05449v2 [math.FA] 13 Dec 2016

FRACTIONAL TYPE MULTILINEAR COMMUTATORS GENERATED BY FRACTIONAL INTEGRAL WITH ROUGH VARIABLE KERNEL AND LOCAL CAMPANATO FUNCTIONS ON GENERALIZED VANISHING LOCAL MORREY SPACES

F.GURBUZ

Abstract. In this paper, we consider the boundedness of fractional type multilinear commutators generated by fractional integral with rough variable kernel and local Campanato functions on both generalized local (central) Morrey spaces and generalized vanishing local Morrey spaces, under generic size conditions which are satisfied by most of the operators in harmonic analysis, respectively.

1. Introduction and main lemmas

The classical Morrey spaces $L_{p,\lambda}$ were introduced by Morrey [23] in 1938 to study the local behavior of solutions of second order elliptic partial differential equations (PDEs). Later, there were many applications of Morrey space to the Navier-Stokes equations (see [21]), the Schrödinger equations (see [29]) and the elliptic problems with discontinuous coefficients (see [4, 12, 25]). The study of the operators of harmonic analysis in vanishing Morrey space, in fact has been almost not touched. A version of the classical Morrey space $L_{p,\lambda}(\mathbb{R}^n)$ where it is possible to approximate by "nice" functions is the so called vanishing Morrey space $VL_{p,\lambda}(\mathbb{R}^n)$ has been introduced by Vitanza in [33] and has been applied there to obtain a regularity result for elliptic PDEs. This is a subspace of functions in $L_{p,\lambda}(\mathbb{R}^n)$, which satisfies the condition

$$\lim_{r\to 0}\sup_{\substack{x\in\mathbb{R}^n\\0<t<r}} t^{-\frac{\lambda}{p}}\|f\|_{L_p(B(x,t))}=0.$$

Later in [34] Vitanza has proved an existence theorem for a Dirichlet problem, under weaker assumptions than in [20] and a $W^{3,2}$ regularity result assuming that the partial derivatives of the coefficients of the highest and lower order terms belong to vanishing Morrey spaces depending on the dimension. Also Ragusa has proved a sufficient condition for commutators of fractional integral operators to belong to vanishing Morrey spaces $VL_{p,\lambda}(\mathbb{R}^n)$ ([27, 28]). For the properties and applications of vanishing Morrey spaces, see also [5].

First of all, we recall some basic definitions and notations used in the paper.

Throughout the paper we assume that $x\in\mathbb{R}^n$ and $r>0$, and also let $B(x,r)$ denote the open ball centered at x of radius r, $B^C(x,r)$ denote its complement and

2010 *Mathematics Subject Classification.* 42B20, 42B25, 42B35.

Key words and phrases. Sublinear operator; fractional integral operator; rough variable kernel; generalized local (central) Morrey spaces; generalized vanishing local Morrey space; fractional type multilinear commutator; local Campanato space.

$|B(x,r)|$ be the Lebesgue measure of the ball $B(x,r)$, and $|B(x,r)| = v_n r^n$, where $v_n = |B(0,1)|$.

Recall that the concept of the generalized local (central) Morrey space $LM_{p,\varphi}^{\{x_0\}}$ has been introduced in [3] and studied in [13, 16].

Definition 1. ***(Generalized local (central) Morrey space)*** *Let $\varphi(x,r)$ be a positive measurable function on $\mathbb{R}^n \times (0,\infty)$ and $1 \le p < \infty$. For any fixed $x_0 \in \mathbb{R}^n$ we denote by $LM_{p,\varphi}^{\{x_0\}} \equiv LM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ the generalized local Morrey space, the space of all functions $f \in L_p^{loc}(\mathbb{R}^n)$ with finite quasinorm*

$$\|f\|_{LM_{p,\varphi}^{\{x_0\}}} = \sup_{r>0} \varphi(x_0,r)^{-1} \left|B(x_0,r)\right|^{-\frac{1}{p}} \|f\|_{L_p(B(x_0,r))} < \infty.$$

According to this definition, we recover the local Morrey space $LL_{p,\lambda}^{\{x_0\}}$ under the choice $\varphi(x_0,r) = r^{\frac{\lambda-n}{p}}$:

$$LL_{p,\lambda}^{\{x_0\}} = LM_{p,\varphi}^{\{x_0\}} \mid_{\varphi(x_0,r)=r^{\frac{\lambda-n}{p}}}.$$

For the properties and applications of generalized local (central) Morrey spaces $LM_{p,\varphi}^{\{x_0\}}$, see also [3, 13, 16].

Definition 2. [3, 13] *Let $1 \le p < \infty$ and $0 \le \lambda < \frac{1}{n}$. A local Campanato function $b \in L_p^{loc}(\mathbb{R}^n)$ is said to belong to the $LC_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$ (local Campanato space), if*

$$\|b\|_{LC_{p,\lambda}^{\{x_0\}}} = \sup_{r>0} \left(\frac{1}{|B(x_0,r)|^{1+\lambda p}} \int\limits_{B(x_0,r)} \left|b(y) - b_{B(x_0,r)}\right|^p dy \right)^{\frac{1}{p}} < \infty, \tag{1.1}$$

where

$$b_{B(x_0,r)} = \frac{1}{|B(x_0,r)|} \int\limits_{B(x_0,r)} b(y)\, dy.$$

Define

$$LC_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n) = \left\{ b \in L_p^{loc}(\mathbb{R}^n) : \|b\|_{LC_{p,\lambda}^{\{x_0\}}} < \infty \right\}.$$

Remark 1. *In* [19]*, Lu and Yang have introduced the central BMO space $CBMO_p(\mathbb{R}^n) = LC_{p,0}^{\{0\}}(\mathbb{R}^n)$. Note that $BMO(\mathbb{R}^n) \subset \bigcap\limits_{p>1} LC_p^{\{x_0\}}(\mathbb{R}^n)$, $1 \le p < \infty$. Moreover, one can imagine that the behavior of $LC_p^{\{x_0\}}(\mathbb{R}^n)$ may be quite different from that of $BMO(\mathbb{R}^n)$ (bounded mean oscillation space), since there is no analogy of the famous John-Nirenberg inequality of $BMO(\mathbb{R}^n)$ for the space $LC_p^{\{x_0\}}(\mathbb{R}^n)$.*

Lemma 1. [3, 13] *Let b be function in $LC_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$, $1 \le p < \infty$, $0 \le \lambda < \frac{1}{n}$ and r_1, $r_2 > 0$. Then*

$$\left(\frac{1}{|B(x_0,r_1)|^{1+\lambda p}} \int\limits_{B(x_0,r_1)} \left|b(y) - b_{B(x_0,r_2)}\right|^p dy \right)^{\frac{1}{p}} \le C \left(1 + \ln \frac{r_1}{r_2}\right) \|b\|_{LC_{p,\lambda}^{\{x_0\}}}, \tag{1.2}$$

where $C > 0$ is independent of b, r_1 and r_2.

From this inequality (1.2), we have

$$\left|b_{B(x_0,r_1)}-b_{B(x_0,r_2)}\right|\leq C\left(1+\ln\frac{r_1}{r_2}\right)\left|B\left(x_0,r_1\right)\right|^{\lambda}\left\|b\right\|_{LC_{p,\lambda}^{\{x_0\}}}, \tag{1.3}$$

and it is easy to see that

$$\left\|b-\left(b\right)_B\right\|_{L_p(B)}\leq C\left(1+\ln\frac{r_1}{r_2}\right)r^{\frac{n}{p}+n\lambda}\left\|b\right\|_{LC_{p,\lambda}^{\{x_0\}}}. \tag{1.4}$$

For brevity, in the sequel we use the following notation

$$\mathfrak{M}_{p,\varphi}\left(f;x_0,r\right):=\frac{\left|B(x_0,r)\right|^{-\frac{1}{p}}\left\|f\right\|_{L_p(B(x_0,r))}}{\varphi(x_0,r)}.$$

Extending the definition of the vanishing Morrey spaces to the case of the generalized local (central) Morrey spaces, we introduce the following definition.

Definition 3. ***(generalized vanishing local Morrey space)*** *The generalized vanishing local Morrey space $VLM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ is defined as the spaces of functions $f\in LM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ such that*

$$\lim_{r\to 0}\mathfrak{M}_{p,\varphi}\left(f;x_0,r\right)=0.$$

Throughout the sequel we assume that

$$\lim_{r\to 0}\frac{1}{\varphi(x_0,r)}=0, \tag{1.5}$$

and

$$\sup_{0<r<\infty}\frac{1}{\varphi(x_0,r)}<\infty, \tag{1.6}$$

which make the spaces $VLM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ non-trivial, since bounded functions with compact support belong to this space. The space $VLM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ is a Banach space with respect to the norm

$$\left\|f\right\|_{VLM_{p,\varphi}^{\{x_0\}}}\equiv\left\|f\right\|_{LM_{p,\varphi}^{\{x_0\}}}=\sup_{r>0}\mathfrak{M}_{p,\varphi}\left(f;x_0,r\right).$$

The space $VLM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ is a closed subspace of the Banach space $LM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$, which may be shown by standard means.

Suppose that S^{n-1} is the unit sphere on $\mathbb{R}^n$ $(n\geq 2)$ equipped with the normalized Lebesgue measure $d\sigma\left(x'\right)$. A function $\Omega\left(x,z\right)$ defined on $\mathbb{R}^n\times\mathbb{R}^n$ is said to belong to $L_\infty(\mathbb{R}^n)\times L_s(S^{n-1})$ for $s>1$, if Ω satisfies the following conditions:

for any x, $z\in\mathbb{R}^n$ and $\lambda>0$,

$$\Omega(x,\lambda z)=\Omega(x,z);$$

$$\left\|\Omega\right\|_{L_\infty(\mathbb{R}^n)\times L_s(S^{n-1})}:=\sup_{x\in\mathbb{R}^n}\left(\int\limits_{S^{n-1}}\left|\Omega(x,z')\right|^s d\sigma\left(z'\right)\right)^{1/s}<\infty,$$

where $z'=z/\left|z\right|$, for any $z\in\mathbb{R}^n\setminus\{0\}$.

Suppose that $T_{\Omega,\alpha}$, $\alpha \in (0,n)$ represents a linear or a sublinear operator with rough variable kernel, which satisfies that for any $f \in S(\mathbb{R}^n)$ with compact support and $x \notin suppf$

$$|T_{\Omega,\alpha} f(x)| \leq c_0 \int_{\mathbb{R}^n} \frac{|\Omega(x,x-y)|}{|x-y|^{n-\alpha}} |f(y)|\, dy, \tag{1.7}$$

where c_0 is independent of f and x. We point out that the condition (1.7) in the case of $\Omega \equiv 1$, $\alpha = 0$ has been introduced by Soria and Weiss in [32]. The condition (1.7) is satisfied by many interesting operators in harmonic analysis, such as the fractional Marcinkiewicz operator, fractional maximal operator, fractional integral operator (Riesz potential) and so on (see [18], [32] for details).

Then, the fractional integral operator with rough variable kernel $\overline{T}_{\Omega,\alpha}$ is defined by

$$\overline{T}_{\Omega,\alpha} f(x) = \int_{\mathbb{R}^n} \frac{\Omega(x,x-y)}{|x-y|^{n-\alpha}} f(y) dy \qquad 0 < \alpha < n,$$

and a related fractional maximal operator with rough variable kernel $M_{\Omega,\alpha}$ is given by

$$M_{\Omega,\alpha} f(x) = \sup_{t>0} |B(x,t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |\Omega(x,x-y)|\, |f(y)| dy \qquad 0 < \alpha < n,$$

where $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$ and these operators also satisfy condition (1.7).

If $\alpha = 0$, then $\overline{T}_{\Omega,\alpha}$ and $M_{\Omega,\alpha}$ becomes the singular integral operator and the Hardy-Littlewood maximal operator by with rough variable kernels, respectively. It is obvious that when $\Omega \equiv 1$, $\overline{T}_{1,\alpha} \equiv \overline{T}_\alpha$ and $M_{1,\alpha} \equiv M_\alpha$ are the fractional integral operator (Riesz potential) and the fractional maximal operator, respectively.

In recent years, the mapping properties of $\overline{T}_{\Omega,\alpha}$ on some kinds of function spaces have been studied in many papers (see [1, 2, 7, 24, 35, 37] for details). In particular, the boundedness of $\overline{T}_{\Omega,\alpha}$ for rough variable kernel in Lebesgue spaces has been obtained by Muckenhoupt and Wheeden [24] as follows:

Lemma 2. [24] *Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. If $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > \frac{n}{n-\alpha}$, for $s' \leq p$ or $q < s$, then we have*

$$\left\| \overline{T}_{\Omega,\alpha} f \right\|_{L_q(\mathbb{R}^n)} \leq C \|\Omega\|_{L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})} \|f\|_{L_p(\mathbb{R}^n)}.$$

Corollary 1. *Under the assumptions of Lemma 2, the operator $M_{\Omega,\alpha}$ from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$ is bounded . Moreover, we have*

$$\|M_{\Omega,\alpha} f\|_{L_q(\mathbb{R}^n)} \leq C \|\Omega\|_{L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})} \|f\|_{L_p(\mathbb{R}^n)}.$$

Proof. Set

$$\widetilde{T}_{|\Omega|,\alpha}(|f|)(x) = \int_{\mathbb{R}^n} \frac{|\Omega(x,x-y)|}{|x-y|^{n-\alpha}} |f(y)|\, dy, \qquad 0 < \alpha < n,$$

where $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})\,(s > 1)$ is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. It is easy to see that, $\widetilde{T}_{|\Omega|,\alpha}$ satisfies Lemma 2. On

the other hand, for any $t>0$, we have

$$\begin{aligned}\widetilde{T}_{|\Omega|,\alpha}\left(|f|\right)(x) &\geq \int\limits_{B(x,t)} \frac{|\Omega(x,x-y)|}{|x-y|^{n-\alpha}}\,|f(y)|\,dy \\ &\geq \frac{1}{t^{n-\alpha}} \int\limits_{B(x,t)} |\Omega(x,x-y)|\,|f(y)|\,dy.\end{aligned}$$

Then by taking supremum for $t>0$, we get

$$M_{\Omega,\alpha}f\left(x\right) \leq C_{n,\alpha}^{-1}\widetilde{T}_{|\Omega|,\alpha}\left(|f|\right)(x) \qquad C_{n,\alpha}=\left|B\left(0,1\right)\right|^{\frac{n-\alpha}{n}}.$$

□

On the other hand, commutators of linear operators take important roles in harmonic analysis and related topics (see [6, 10, 11, 13, 15, 16, 17, 26, 30]). There are two major reasons for considering the problem of commutators. The first one is that the boundedness of commutators can produce some characterizations of function spaces (see [3, 6, 13, 14, 15, 16, 17, 26, 30]). The other one is that the theory of commutators plays an important role in the study of the regularity of solutions to elliptic and parabolic PDEs of the second order (see [8, 9, 12, 31]).

Let $b_i\,(i=1,\ldots,m)$ be locally integrable functions on $\mathbb{R}^n$, then the multilinear commutators generated by fractional integral and maximal operator with rough variable kernel and $\overrightarrow{b}=(b_1,\ldots,b_m)$ are given as follows, respectively:

$$[\overrightarrow{b},\overline{T}_{\Omega,\alpha}]f\left(x\right)=\int\limits_{\mathbb{R}^n}\prod_{i=1}^{m}\left[b_i\left(x\right)-b_i\left(y\right)\right]\frac{\Omega(x,x-y)}{|x-y|^{n-\alpha}}f\left(y\right)dy, \qquad 0<\alpha<n,$$

$$M_{\Omega,\overrightarrow{b},\alpha}f\left(x\right)=\sup_{t>0}|B(x,t)|^{-1+\frac{\alpha}{n}}\int\limits_{B(x,t)}\prod_{i=1}^{m}\left[b_i\left(x\right)-b_i\left(y\right)\right]|\Omega(x,x-y)|\,|f(y)|dy, \qquad 0<\alpha<n.$$

For $m=1$, $[\overrightarrow{b},\overline{T}_{\Omega,\alpha}]$ and $M_{\Omega,\overrightarrow{b},\alpha}$ are obviously the commutator operators of $\overline{T}_{\Omega,\alpha}$ and $M_{\Omega,\alpha}$,

$$\begin{aligned}[b,\overline{T}_{\Omega,\alpha}]f(x) &\equiv b(x)\overline{T}_{\Omega,\alpha}f(x)-\overline{T}_{\Omega,\alpha}(bf)(x) \\ &=\int\limits_{\mathbb{R}^n}[b(x)-b(y)]\frac{\Omega(x,x-y)}{|x-y|^{n-\alpha}}f(y)dy,\end{aligned}$$

and

$$\begin{aligned}M_{\Omega,b,\alpha}\left(f\right)(x) &\equiv b\left(x\right)M_{\Omega,\alpha}f\left(x\right)-M_{\Omega,\alpha}\left(bf\right)(x) \\ &=\sup_{t>0}|B(x,t)|^{-1+\frac{\alpha}{n}}\int\limits_{B(x,t)}|b\left(x\right)-b\left(y\right)|\,|\Omega(x,x-y)|\,|f(y)|dy,\end{aligned}$$

where $0<\alpha<n$ and f is a suitable function.

In [22], the authors obtain the boundedness for the multilinear commutators generated by singular integral operators with rough variable kernel and local Campanato functions on generalized local Morrey spaces.

Inspired by [22], in this paper we give local Campanato space estimates for fractional type multilinear commutators with rough variable kernel on both generalized local Morrey spaces and generalized vanishing local Morrey spaces, respectively.

But, the techniques and non-trivial estimates which have been used in the proofs of our main results are quite different from [22]. For example, using inequality about the weighted Hardy operator H_w in [22], in this paper we will only use the following relationship between essential supremum and essential infimum

$$\left(\operatorname*{essinf}_{x\in E} f(x)\right)^{-1} = \operatorname*{esssup}_{x\in E} \frac{1}{f(x)}, \tag{1.8}$$

where f is any real-valued nonnegative function and measurable on E (see [36], page 143).

We first need some lemmas (our main lemmas) which are used in the proof of the main results. These lemmas with their proofs can be formulated as follows, respectively:

Lemma 3. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. Let $T_{\Omega,\alpha}$ be a sublinear operator satisfying condition (1.7), bounded from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$.*

If $p > 1$ and $s' \le p$, then the inequality

$$\|T_{\Omega,\alpha} f\|_{L_q(B(x_0,r))} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))}\, dt \tag{1.9}$$

holds for any ball $B(x_0, r)$ and for all $f \in L_p^{loc}(\mathbb{R}^n)$.

If $p > 1$ and $q < s$, then the inequality

$$\|T_{\Omega,\alpha} f\|_{L_q(B(x_0,r))} \lesssim r^{\frac{n}{q}-\frac{n}{s}} \int_{2r}^{\infty} t^{\frac{n}{s}-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))}\, dt$$

holds for any ball $B(x_0, r)$ and for all $f \in L_p^{loc}(\mathbb{R}^n)$.

Proof. Let $0 < \alpha < n$, $1 \le s' < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. Set $B = B(x_0, r)$ for the ball centered at x_0 and of radius r and $2B = B(x_0, 2r)$. We represent f as

$$f = f_1 + f_2, \qquad f_1(y) = f(y)\chi_{2B}(y), \qquad f_2(y) = f(y)\chi_{(2B)^C}(y), \qquad r > 0 \tag{1.10}$$

and have

$$\|T_{\Omega,\alpha} f\|_{L_q(B)} \le \|T_{\Omega,\alpha} f_1\|_{L_q(B)} + \|T_{\Omega,\alpha} f_2\|_{L_q(B)}.$$

Since $f_1 \in L_p(\mathbb{R}^n)$, $T_{\Omega,\alpha} f_1 \in L_q(\mathbb{R}^n)$ and by the boundedness of $T_{\Omega,\alpha}$ from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$ (see Lemma 2) it follows that:

$$\|T_{\Omega,\alpha} f_1\|_{L_q(B)} \le \|T_{\Omega,\alpha} f_1\|_{L_q(\mathbb{R}^n)} \le C \|f_1\|_{L_p(\mathbb{R}^n)} = C \|f\|_{L_p(2B)},$$

where constant $C > 0$ is independent of f.

It is clear that $x \in B$, $y \in (2B)^C$ implies $\frac{1}{2}|x_0 - y| \le |x - y| \le \frac{3}{2}|x_0 - y|$. Then we get

$$|T_{\Omega,\alpha} f_2(x)| \le 2^{n-\alpha} c_1 \int_{(2B)^C} \frac{|f(y)|\,|\Omega(x,\ x-y)|}{|x_0 - y|^{n-\alpha}} dy.$$

By Fubini's theorem, we have

$$\begin{aligned}\int\limits_{(2B)^C}\frac{|f(y)|\,|\Omega(x,x-y)|}{|x_0-y|^{n-\alpha}}dy &\approx \int\limits_{(2B)^C}|f(y)|\,|\Omega(x,x-y)|\int\limits_{|x_0-y|}^{\infty}\frac{dt}{t^{n+1-\alpha}}dy\\ &\approx \int\limits_{2r}^{\infty}\int\limits_{2r\le|x_0-y|\le t}|f(y)|\,|\Omega(x,x-y)|\,dy\frac{dt}{t^{n+1-\alpha}}\\ &\lesssim \int\limits_{2r}^{\infty}\int\limits_{B(x_0,t)}|f(y)|\,|\Omega(x,x-y)|\,dy\frac{dt}{t^{n+1-\alpha}}.\end{aligned}$$

Applying Hölder's inequality, we get

$$\begin{aligned}&\int\limits_{(2B)^C}\frac{|f(y)|\,|\Omega(x,x-y)|}{|x_0-y|^{n-\alpha}}dy\\ &\lesssim \int\limits_{2r}^{\infty}\|f\|_{L_p(B(x_0,t))}\,\|\Omega(x,x-\cdot)\|_{L_s(B(x_0,t))}\,|B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{s}}\,\frac{dt}{t^{n+1-\alpha}}.\end{aligned}\tag{1.11}$$

For $x\in B(x_0,t)$, notice that Ω is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$ and $\Omega\in L_\infty(\mathbb{R}^n)\times L_s(S^{n-1})$, $s>1$. Then, we obtain

$$\begin{aligned}\left(\int\limits_{B(x_0,t)}|\Omega(x,x-y)|^s\,dy\right)^{\frac{1}{s}} &= \left(\int\limits_{B(x-x_0,t)}|\Omega(x,z)|^s\,dz\right)^{\frac{1}{s}}\\ &\le \left(\int\limits_{B(0,t+|x-x_0|)}|\Omega(x,z)|^s\,dz\right)^{\frac{1}{s}}\\ &\le \left(\int\limits_{B(0,2t)}|\Omega(x,z)|^s\,dz\right)^{\frac{1}{s}}\\ &= \left(\int\limits_{S^{n-1}}\int\limits_0^{2t}|\Omega(x,z')|^s\,d\sigma(z')\,r^{n-1}dr\right)^{\frac{1}{s}}\\ &= C\,\|\Omega\|_{L_\infty(\mathbb{R}^n)\times L_s(S^{n-1})}\,|B(x_0,2t)|^{\frac{1}{s}}.\end{aligned}\tag{1.12}$$

Thus, by (1.12), it follows that:

$$|T_{\Omega,\alpha}f_2(x)|\lesssim\int\limits_{2r}^{\infty}\|f\|_{L_p(B(x_0,t))}\,\frac{dt}{t^{\frac{n}{q}+1}}.$$

Moreover, for all $p\in(1,\infty)$ the inequality

$$\|T_{\Omega,\alpha}f_2\|_{L_q(B)}\lesssim r^{\frac{n}{q}}\int\limits_{2r}^{\infty}\|f\|_{L_p(B(x_0,t))}\,\frac{dt}{t^{\frac{n}{q}+1}}\tag{1.13}$$

is valid. Thus, we obtain

$$\|T_{\Omega,\alpha}f\|_{L_q(B)} \lesssim \|f\|_{L_p(2B)} + r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

On the other hand, we have

$$\begin{aligned} \|f\|_{L_p(2B)} &\approx r^{\frac{n}{q}} \|f\|_{L_p(2B)} \int_{2r}^{\infty} \frac{dt}{t^{\frac{n}{q}+1}} \\ &\leq r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}}. \end{aligned} \tag{1.14}$$

By combining the above inequalities, we obtain

$$\|T_{\Omega,\alpha}f\|_{L_q(B)} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))}\, dt.$$

For the case of $1 < q < s$, we can also use the same method, so we omit the details. This completes the proof of Lemma 3. □

Lemma 4. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $T_{\Omega,\alpha}$ be a linear operator satisfying condition (1.7). Let also $0 < \alpha < n$ and $1 < q, q_1, p_i, p < \frac{n}{\alpha}$ with $\frac{1}{q} = \sum_{i=1}^{m} \frac{1}{p_i} + \frac{1}{p}$, $\frac{1}{q_1} = \frac{1}{q} - \frac{\alpha}{n}$ and $\overrightarrow{b} \in LC^{\{x_0\}}_{p_i,\lambda_i}(\mathbb{R}^n)$ for $0 \leq \lambda_i < \frac{1}{n}$, $i = 1, \ldots, m$. Then, for $s' \leq q$ the inequality*

(1.15)

$$\|[\overrightarrow{b}, T_{\Omega,\alpha}]f\|_{L_{q_1}(B(x_0,r))} \lesssim \prod_{i=1}^{m} \|\overrightarrow{b}\|_{LC^{\{x_0\}}_{p_i,\lambda_i}} r^{\frac{n}{q_1}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1} + \left(\sum_{i=1}^{m} \lambda_i + \sum_{i=1}^{m} \frac{1}{p_i}\right)\right)-1} \|f\|_{L_p(B(x_0,t))} dt$$

holds for any ball $B(x_0, r)$ and for all $f \in L_p^{loc}(\mathbb{R}^n)$. Also, for $q_1 < s$ the inequality

$$\|[\overrightarrow{b}, T_{\Omega,\alpha}]f\|_{L_{q_1}(B(x_0,r))} \lesssim \prod_{i=1}^{m} \|\overrightarrow{b}\|_{LC^{\{x_0\}}_{p_i,\lambda_i}} r^{\frac{n}{q_1}-\frac{n}{s}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1} + \left(\frac{1}{s} + \sum_{i=1}^{m} \lambda_i + \sum_{i=1}^{m} \frac{1}{p_i}\right)\right)-1} \|f\|_{L_p(B(x_0,t))} dt$$

holds for any ball $B(x_0, r)$ and for all $f \in L_p^{loc}(\mathbb{R}^n)$.

Proof. Without loss of generality, it is sufficient to show that the conclusion holds for $[\overrightarrow{b}, T_{\Omega,\alpha}]f = [(b_1, b_2), T_{\Omega,\alpha}]f$. As in the proof of Lemma 3, we represent f in form (1.10) and thus have

$$\|[(b_1, b_2), T_{\Omega,\alpha}]f\|_{L_{q_1}(B)} \leq \|[(b_1, b_2), T_{\Omega,\alpha}]f_1\|_{L_{q_1}(B)} + \|[(b_1, b_2), T_{\Omega,\alpha}]f_2\|_{L_{q_1}(B)} =: F + G.$$

Let us estimate $F + G$, respectively.

For $[(b_1,b_2),T_{\Omega,\alpha}]f_1(x)$, we have the following decomposition,

$$\begin{aligned}[(b_1,b_2),T_{\Omega,\alpha}]f_1(x) &= (b_1(x)-(b_1)_B)(b_2(x)-(b_2)_B)T_{\Omega,\alpha}f_1(x)\\ &\quad-(b_1(\cdot)-(b_1)_B)T_{\Omega,\alpha}((b_2(\cdot)-(b_2)_B)f_1)(x)\\ &\quad+(b_2(x)-(b_2)_B)T_{\Omega,\alpha}((b_1(x)-(b_1)_B)f_1)(x)-\\ &\quad T_{\Omega,\alpha}((b_1(\cdot)-(b_1)_B)(b_2(\cdot)-(b_2)_B)f_1)(x).\end{aligned}$$

Hence, we get

$$\begin{aligned}F &= \|[(b_1,b_2),T_{\Omega,\alpha}]f_1\|_{L_{q_1}(B)} \lesssim\\ &\quad\|(b_1-(b_1)_B)(b_2(x)-(b_2)_B)T_{\Omega,\alpha}f_1\|_{L_{q_1}(B)}\\ &\quad+\|(b_1-(b_1)_B)T_{\Omega,\alpha}((b_2-(b_2)_B)f_1)\|_{L_{q_1}(B)}\\ &\quad+\|(b_2-(b_2)_B)T_{\Omega,\alpha}((b_1-(b_1)_B)f_1)\|_{L_{q_1}(B)}\\ &\quad+\|T_{\Omega,\alpha}((b_1-(b_1)_B)(b_2-(b_2)_B)f_1)\|_{L_{q_1}(B)}\\ &\equiv F_1+F_2+F_3+F_4. \qquad (1.16)\end{aligned}$$

One observes that the estimate of F_2 is analogous to that of F_3. Thus, we will only estimate F_1, F_2 and F_4.

To estimate F_1, let $1<\overline{q},\overline{r}<\infty$, such that$\frac{1}{\overline{q}}=\frac{1}{p}-\frac{\alpha}{n}$, $\frac{1}{q_1}=\frac{1}{\overline{r}}+\frac{1}{\overline{q}}$,$\frac{1}{\overline{r}}=\frac{1}{p_1}+\frac{1}{p_2}$. Then, using Hölder's inequality and by the boundedness of $T_{\Omega,\alpha}$ from L_p into $L_{\overline{q}}$ (see Lemma 2) it follows that:

$$\begin{aligned}F_1 &= \|(b_1-(b_1)_B)(b_2(x)-(b_2)_B)T_{\Omega,\alpha}f_1\|_{L_{q_1}(B)}\\ &\lesssim \|(b_1-(b_1)_B)(b_2-(b_2)_B)\|_{L_{\overline{r}}(B)}\|T_{\Omega,\alpha}f_1\|_{L_{\overline{q}}(B)}\\ &\lesssim \|b_1-(b_1)_B\|_{L_{p_1}(B)}\|b_2-(b_2)_B\|_{L_{p_2}(B)}\|f\|_{L_p(2B)}\\ &\lesssim \|b_1-(b_1)_B\|_{L_{p_1}(B)}\|b_2-(b_2)_B\|_{L_{p_2}(B)}r^{n\left(\frac{1}{p}-\frac{\alpha}{n}\right)}\int_{2r}^{\infty}\|f\|_{L_{q_1}(B(x_0,t))}\frac{dt}{t^{\frac{n}{p}+1-\alpha}}.\end{aligned}$$

From Lemma 1, it is easy to see that

$$\|b_i-(b_i)_B\|_{L_{p_i}(B)}\le Cr^{\frac{n}{p_i}+n\lambda_i}\|b_i\|_{LC^{\{x_0\}}_{p_i,\lambda_i}}, \qquad (1.17)$$

and

$$\begin{aligned}\|b_i-(b_i)_B\|_{L_{p_i}(2B)} &\le \|b_i-(b_i)_{2B}\|_{L_{p_i}(2B)}+\|(b_i)_B-(b_i)_{2B}\|_{L_{p_i}(2B)}\\ &\lesssim r^{\frac{n}{p_i}+n\lambda_i}\|b_i\|_{LC^{\{x_0\}}_{p_i,\lambda_i}}, \qquad (1.18)\end{aligned}$$

for $i=1,2$. Hence, by (1.17) we get

$$\begin{aligned}F_1 &\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{n\left(\frac{1}{p_1}+\frac{1}{p_2}+\frac{1}{p}-\frac{\alpha}{n}\right)}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{p}+n(\lambda_1+\lambda_2)-1+\alpha}\|f\|_{L_p(B(x_0,t))}\,dt\\ &\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.\end{aligned}$$

To estimate F_2, let $1<\tau<\infty$, such that $\frac{1}{q_1}=\frac{1}{p_1}+\frac{1}{\tau}$. Then, similar to the estimates for F_1, we have

$$\begin{aligned}F_2&=\left\|\left(b_1-(b_1)_B\right)T_{\Omega,\alpha}\left(\left(b_2-(b_2)_B\right)f_1\right)\right\|_{L_{q_1}(B)}\\&\lesssim\left\|b_1-(b_1)_B\right\|_{L_{p_1}(B)}\left\|T_{\Omega,\alpha}\left(\left(b_2\left(\cdot\right)-(b_2)_B\right)f_1\right)\right\|_{L_\tau(B)}\\&\lesssim\left\|b_1-(b_1)_B\right\|_{L_{p_1}(B)}\left\|\left(b_2\left(\cdot\right)-(b_2)_B\right)f_1\right\|_{L_k(B)}\\&\lesssim\left\|b_1-(b_1)_B\right\|_{L_{p_1}(B)}\left\|b_2-(b_2)_B\right\|_{L_{p_2}(2B)}\left\|f\right\|_{L_p(2B)},\end{aligned}$$

where $1<k<\frac{2n}{\alpha}$, such that $\frac{1}{k}=\frac{1}{p_2}+\frac{1}{p}=\frac{1}{\tau}+\frac{\alpha}{n}$. By (1.17) and (1.18), we get

$$F_2\lesssim\left\|b_1\right\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\left\|b_2\right\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\left\|f\right\|_{L_p(B(x_0,t))}dt.$$

In a similar way, F_3 has the same estimate as above, so we omit the details. Then we have that

$$F_3\lesssim\left\|b_1\right\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\left\|b_2\right\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\left\|f\right\|_{L_p(B(x_0,t))}dt.$$

Now let us consider the term F_4. Let $1<q,\tau<\frac{2n}{\alpha}$, such that $\frac{1}{q}=\frac{1}{\tau}+\frac{1}{p}$, $\frac{1}{\tau}=\frac{1}{p_1}+\frac{1}{p_2}$ and $\frac{1}{q_1}=\frac{1}{q}-\frac{\alpha}{n}$. Then by the boundedness of $T_{\Omega,\alpha}$ from L_q into L_{q_1} (see Lemma 2), Hölder's inequality and (1.18), we obtain

$$\begin{aligned}F_4&=\left\|T_{\Omega,\alpha}\left(\left(b_1-(b_1)_B\right)\left(b_2-(b_2)_B\right)f_1\right)\right\|_{L_{q_1}(B)}\\&\lesssim\left\|\left(b_1-(b_1)_B\right)\left(b_2-(b_2)_B\right)f_1\right\|_{L_q(B)}\\&\lesssim\left\|\left(b_1-(b_1)_B\right)\left(b_2-(b_2)_B\right)\right\|_{L_\tau(B)}\left\|f_1\right\|_{L_p(B)}\\&\lesssim\left\|b_1-(b_1)_B\right\|_{L_{p_1}(2B)}\left\|b_2-(b_2)_B\right\|_{L_{p_2}(2B)}\left\|f\right\|_{L_p(2B)}\\&\lesssim\left\|b_1\right\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\left\|b_2\right\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\left\|f\right\|_{L_p(B(x_0,t))}dt.\end{aligned}$$

Combining all the estimates of F_1, F_2, F_3, F_4; we get

$$\begin{aligned}F=\left\|\left[\left(b_1,b_2\right),T_{\Omega,\alpha}\right]f_1\right\|_{L_{q_1}(B)}&\lesssim\left\|b_1\right\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\left\|b_2\right\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}\\&\times\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\left\|f\right\|_{L_p(B(x_0,t))}dt.\end{aligned}$$

Now, let us estimate $G = \|[(b_1, b_2), T_{\Omega,\alpha}]f_2\|_{L_{q_1}(B)}$. For G, it's similar to (1.16) we also write

$$\begin{aligned} G &= \|[(b_1, b_2), T_{\Omega,\alpha}]f_2\|_{L_{q_1}(B)} \lesssim \\ &\|(b_1 - (b_1)_B)(b_2(x) - (b_2)_B) T_{\Omega,\alpha} f_2\|_{L_{q_1}(B)} \\ &+ \|(b_1 - (b_1)_B) T_{\Omega,\alpha}((b_2 - (b_2)_B) f_2)\|_{L_{q_1}(B)} \\ &+ \|(b_2 - (b_2)_B) T_{\Omega,\alpha}((b_1 - (b_1)_B) f_2)\|_{L_{q_1}(B)} \\ &+ \|T_{\Omega,\alpha}((b_1 - (b_1)_B)(b_2 - (b_2)_B) f_2)\|_{L_{q_1}(B)} \\ &\equiv G_1 + G_2 + G_3 + G_4. \end{aligned}$$

To estimate G_1, let $1 < p_1, p_2 < \frac{2n}{\alpha}$, such that $\frac{1}{q_1} = \frac{1}{\overline{p}} + \frac{1}{\overline{q}}$, $\frac{1}{\overline{p}} = \frac{1}{p_1} + \frac{1}{p_2}$ and $\frac{1}{\overline{q}} = \frac{1}{p} - \frac{\alpha}{n}$. Then, using Hölder's inequality, noting that in (1.13) and by (1.17), we have

$$\begin{aligned} G_1 &= \|(b_1 - (b_1)_B)(b_2(x) - (b_2)_B) T_{\Omega,\alpha} f_2\|_{L_{q_1}(B)} \\ &\lesssim \|(b_1 - (b_1)_B)(b_2 - (b_2)_B)\|_{L_{\overline{p}}(B)} \|T_{\Omega,\alpha} f_2\|_{L_{\overline{q}}(B)} \\ &\lesssim \|b_1 - (b_1)_B\|_{L_{p_1}(B)} \|b_2 - (b_2)_B\|_{L_{p_2}(B)} r^{\frac{n}{\overline{q}}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} t^{-\frac{n}{\overline{q}}-1} dt \\ &\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{n\left(\frac{1}{p_1}+\frac{1}{p_2}+\frac{1}{p}-\frac{\alpha}{n}\right)} \\ &\times \int_{2r}^{\infty} \left(1 + \ln\frac{t}{r}\right)^2 t^{-\frac{n}{p}+n(\lambda_1+\lambda_2)-1+\alpha} \|f\|_{L_p(B(x_0,t))} dt \\ &\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{q_1}} \int_{2r}^{\infty} \left(1 + \ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1} \|f\|_{L_p(B(x_0,t))} dt. \end{aligned}$$

On the other hand, for the estimates used in G_2, G_3, we have to prove the below inequality:

(1.19)

$$|T_{\Omega,\alpha}((b_2 - (b_2)_B) f_2)(x)| \lesssim \|b\|_{LC^{\{x_0\}}_{p_2,\lambda}} \int_{2r}^{\infty} \left(1 + \ln\frac{t}{r}\right) t^{-\frac{n}{p}+n\lambda_2-1+\alpha} \|f\|_{L_p(B(x_0,t))} dt.$$

Indeed, when $s' \le q$, for $x \in B$, by Fubini's theorem and applying Hölder's inequality and from (1.3), (1.4), (1.12) we have

$$\begin{aligned} |T_{\Omega,\alpha}((b_2 - (b_2)_B) f_2)(x)| &\lesssim \int_{(2B)^C} |b_2(y) - (b_2)_B| |\Omega(x, x-y)| \frac{|f(y)|}{|x-y|^{n-\alpha}} dy \\ &\lesssim \int_{(2B)^C} |b_2(y) - (b_2)_B| |\Omega(x, x-y)| \frac{|f(y)|}{|x_0-y|^{n-\alpha}} dy \\ &\approx \int_{2r}^{\infty} \int_{2r<|x_0-y|<t} |b_2(y) - (b_2)_B| |\Omega(x, x-y)| |f(y)| dy \frac{dt}{t^{n-\alpha+1}} \end{aligned}$$

$$\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} \left| b_2(y) - (b_2)_{B(x_0,t)} \right| |\Omega(x, x-y)| \, |f(y)| \, dy \frac{dt}{t^{n-\alpha+1}}$$
$$+ \int_{2r}^{\infty} \left| (b_2)_{B(x_0,r)} - (b_2)_{B(x_0,t)} \right| \int_{B(x_0,t)} |\Omega(x, x-y)| \, |f(y)| \, dy \frac{dt}{t^{n-\alpha+1}}$$
$$\lesssim \int_{2r}^{\infty} \left\| b_2(\cdot) - (b_2)_{B(x_0,t)} \right\|_{L_{p_2}(B(x_0,t))} \|\Omega(x, x-\cdot)\|_{L_s(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p_2}-\frac{1}{s}-\frac{1}{p}} \frac{dt}{t^{n-\alpha+1}}$$
$$+ \int_{2r}^{\infty} \left| (b_2)_{B(x_0,r)} - (b_2)_{B(x_0,t)} \right| \|f\|_{L_p(B(x_0,t))} \|\Omega(x, x-\cdot)\|_{L_s(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{s}} \frac{dt}{t^{n-\alpha+1}}$$
$$\lesssim \int_{2r}^{\infty} \left\| b_2(\cdot) - (b_2)_{B(x_0,t)} \right\|_{L_{p_2}(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n\left(\frac{1}{p_2}+\frac{1}{p}\right)-\alpha+1}}$$
$$+ \|b\|_{LC^{\{x_0\}}_{p_2,\lambda}} \int_{2r}^{\infty} \left(1 + \ln \tfrac{t}{r}\right) t^{-\frac{n}{p}+n\lambda_2-1+\alpha} \|f\|_{L_p(B(x_0,t))} \, dt$$
$$\lesssim \|b\|_{LC^{\{x_0\}}_{p_2,\lambda}} \int_{2r}^{\infty} \left(1 + \ln \tfrac{t}{r}\right) t^{-\frac{n}{p}+n\lambda_2-1+\alpha} \|f\|_{L_p(B(x_0,t))} \, dt.$$

This completes the proof of inequality (1.19).

Let $1 < \tau < \infty$, such that $\frac{1}{q_1} = \frac{1}{p_1} + \frac{1}{\tau}$ and $\frac{1}{\tau} = \frac{1}{p_2} + \frac{1}{p} - \frac{\alpha}{n}$. Then, using Hölder's inequality and from (1.19) and (1.4), we get

$$G_2 = \|(b_1 - (b_1)_B) T_{\Omega,\alpha}((b_2 - (b_2)_B) f_2)\|_{L_{q_1}(B)}$$
$$\lesssim \|b_1 - (b_1)_B\|_{L_{p_1}(B)} \|T_{\Omega,\alpha}((b_2(\cdot) - (b_2)_B) f_2)\|_{L_\tau(B)}$$
$$\lesssim \|b_1 - (b_1)_B\|_{L_{p_1}(B)} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{\tau}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{p}+n\lambda_2-1+\alpha} \|f\|_{L_p(B(x_0,t))} \, dt$$
$$\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{q_1}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1} \|f\|_{L_p(B(x_0,t))} \, dt.$$

Similarly, G_3 has the same estimate above, so here we omit the details. Then the inequality

$$G_3 = \|(b_2 - (b_2)_B) T_{\Omega,\alpha}((b_1 - (b_1)_B) f_2)\|_{L_{q_1}(B)}$$
$$\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{q_1}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1} \|f\|_{L_p(B(x_0,t))} \, dt$$

is valid.

Now, let us estimate $G_4 = \|T_{\Omega,\alpha}((b_1 - (b_1)_B)(b_2 - (b_2)_B) f_2)\|_{L_{q_1}(B)}$. It's similar to the estimate of (1.19), for any $x \in B$, we also write

$|T_{\Omega,\alpha}((b_1 - (b_1)_B)(b_2 - (b_2)_B) f_2)(x)|$

$$\lesssim \int_{2r}^{\infty}\int_{B(x_0,t)} \left|b_1(y)-(b_1)_{B(x_0,t)}\right|\left|b_2(y)-(b_2)_{B(x_0,t)}\right| |\Omega(x,x-y)|\,|f(y)|\,dy\tfrac{dt}{t^{n-\alpha+1}}$$
$$+\int_{2r}^{\infty}\int_{B(x_0,t)} \left|b_1(y)-(b_1)_{B(x_0,t)}\right|\left|(b_2)_{B(x_0,t)}-(b_2)_{B(x_0,r)}\right| |\Omega(x,x-y)|\,|f(y)|\,dy\tfrac{dt}{t^{n-\alpha+1}}$$
$$+\int_{2r}^{\infty}\int_{B(x_0,t)} \left|(b_1)_{B(x_0,t)}-(b_2)_{B(x_0,r)}\right|\left|b_2(y)-(b_2)_{B(x_0,t)}\right| |\Omega(x,x-y)|\,|f(y)|\,dy\tfrac{dt}{t^{n-\alpha+1}}$$
$$+\int_{2r}^{\infty}\int_{B(x_0,t)} \left|(b_1)_{B(x_0,t)}-(b_2)_{B(x_0,r)}\right|\left|(b_2)_{B(x_0,t)}-(b_2)_{B(x_0,r)}\right| |\Omega(x,x-y)|\,|f(y)|\,dy\tfrac{dt}{t^{n-\alpha+1}}$$
$$\equiv G_{41}+G_{42}+G_{43}+G_{44}.$$

Let us estimate G_{41}, G_{42}, G_{43}, G_{44}, respectively.

Firstly, to estimate G_{41}, similar to the estimate of (1.19), we get

$$G_{41}\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.$$

Secondly, to estimate G_{42} and G_{43}, from (1.19), (1.3) and (1.4), it follows that

$$G_{42}\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt,$$

and

$$G_{43}\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.$$

Finally, to estimate G_{44}, similar to the estimate of (1.19) and from (1.3) and (1.4), we have

$$G_{44}\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.$$

By the estimates of G_{4j} above, where $j=1,\,2,\,3$, we know that

$$|T_{\Omega,\alpha}\left((b_1-(b_1)_B)(b_2-(b_2)_B)f_2\right)(x)|\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}$$
$$\times\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.$$

Then, we have

$$G_4=\|T_{\Omega,\alpha}\left((b_1-(b_1)_B)(b_2-(b_2)_B)f_2\right)\|_{L_{q_1}(B)}\lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}}\|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}}r^{\frac{n}{q_1}}$$
$$\times\int_{2r}^{\infty}\left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1}\|f\|_{L_p(B(x_0,t))}\,dt.$$

So, combining all the estimates for G_1, G_2, G_3, G_4, we get

$$G = \|[(b_1,b_2),T_{\Omega,\alpha}]f_2\|_{L_{q_1}(B)} \lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{q_1}} \times \int_{2r}^{\infty} \left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1} \|f\|_{L_p(B(x_0,t))}\,dt.$$

Thus, putting estimates F and G together, we get the desired conclusion

$$\|[(b_1,b_2),T_{\Omega,\alpha}]f\|_{L_{q_1}(B(x_0,r))} \lesssim \|b_1\|_{LC^{\{x_0\}}_{p_1,\lambda_1}} \|b_2\|_{LC^{\{x_0\}}_{p_2,\lambda_2}} r^{\frac{n}{q_1}} \times \int_{2r}^{\infty} \left(1+\ln\frac{t}{r}\right)^2 t^{-\frac{n}{q_1}+n(\lambda_1+\lambda_2)+n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)-1} \|f\|_{L_p(B(x_0,t))}\,dt.$$

For the case of $q_1 < s$, we can also use the same method, so we omit the details. This completes the proof of Lemma 4. □

At last, throughout the paper we use the letter C for a positive constant, independent of appropriate parameters and not necessarily the same at each occurrence. By $A \lesssim B$ we mean that $A \le CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2. Main Results

Now we are ready to give the following main results with their proofs , respectively.

Theorem 1. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $T_{\Omega,\alpha}$ be a linear operator satisfying condition (1.7). Let also $0 < \alpha < n$ and $1 < q, q_1, p_i, p < \frac{n}{\alpha}$ with $\frac{1}{q} = \sum_{i=1}^{m} \frac{1}{p_i} + \frac{1}{p}$, $\frac{1}{q_1} = \frac{1}{q} - \frac{\alpha}{n}$ and $\overrightarrow{b} \in LC^{\{x_0\}}_{p_i,\lambda_i}(\mathbb{R}^n)$ for $0 \le \lambda_i < \frac{1}{n}$, $i = 1,\dots,m$.*

Let also, for $s' \le q$ the pair (φ_1,φ_2) satisfies the condition

$$\int_r^{\infty} \left(1+\ln\frac{t}{r}\right)^m \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1}-\left(\sum_{i=1}^{m}\lambda_i+\sum_{i=1}^{m}\frac{1}{p_i}\right)\right)+1}} \le C\,\varphi_2(x_0,r), \tag{2.1}$$

and for $q_1 < s$ the pair (φ_1,φ_2) satisfies the condition

$$\int_r^{\infty} \left(1+\ln\frac{t}{r}\right)^m \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1}-\left(\frac{1}{s}+\sum_{i=1}^{m}\lambda_i+\sum_{i=1}^{m}\frac{1}{p_i}\right)\right)+1}}\,dt \le C\,\varphi_2(x_0,r) r^{\frac{n}{s}}, \tag{2.2}$$

where C does not depend on r.

Then, the operator $[\overrightarrow{b},T_{\Omega,\alpha}]$ is bounded from $LM^{\{x_0\}}_{p,\varphi_1}$ to $LM^{\{x_0\}}_{q_1,\varphi_2}$. Moreover,

$$\left\|[\overrightarrow{b},T_{\Omega,\alpha}]\right\|_{LM^{\{x_0\}}_{q_1,\varphi_2}} \lesssim \prod_{i=1}^{m} \|\overrightarrow{b}\|_{LC^{\{x_0\}}_{p_i,\lambda_i}} \|f\|_{LM^{\{x_0\}}_{p,\varphi_1}}.$$

Proof. Since $f \in LM_{p_1,\varphi_1}^{\{x_0\}}$, by (1.8) and it is also non-decreasing, with respect to t, of the norm $\|f\|_{L_{p_1}(B(x_0,t))}$, we get

$$\frac{\|f\|_{L_p(B(x_0,t))}}{\operatorname*{essinf}_{0<t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}} \leq \operatorname*{esssup}_{0<t<\tau<\infty} \frac{\|f\|_{L_p(B(x_0,t))}}{\varphi_1(x_0,\tau)\tau^{\frac{n}{p}}}$$

$$\leq \operatorname*{esssup}_{0<\tau<\infty} \frac{\|f\|_{L_p(B(x_0,\tau))}}{\varphi_1(x_0,\tau)\tau^{\frac{n}{p}}} \leq \|f\|_{LM_{p,\varphi_1}^{\{x_0\}}}. \tag{2.3}$$

For $s' \leq q < \infty$, since (φ_1, φ_2) satisfies (2.1) and by (2.3), we have

$$\int_r^\infty \left(1+\ln\frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1}+\left(\sum_{i=1}^m \lambda_i+\sum_{i=1}^m \frac{1}{p_i}\right)\right)-1} \|f\|_{L_p(B(x_0,t))}dt$$

$$\leq \int_r^\infty \left(1+\ln\frac{t}{r}\right)^m \frac{\|f\|_{L_p(B(x_0,t))}}{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}} \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1}-\left(\sum_{i=1}^m \lambda_i+\sum_{i=1}^m \frac{1}{p_i}\right)\right)+1}}dt$$

$$\leq C\|f\|_{LM_{p,\varphi_1}^{\{x_0\}}} \int_r^\infty \left(1+\ln\frac{t}{r}\right)^m \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1}-\left(\sum_{i=1}^m \lambda_i+\sum_{i=1}^m \frac{1}{p_i}\right)\right)+1}}dt$$

$$\leq C\|f\|_{LM_{p,\varphi_1}^{\{x_0\}}} \varphi_2(x_0,r). \tag{2.4}$$

Then by (1.15) and (2.4), we get

$$\begin{aligned}
\left\|[\overrightarrow{b}, T_{\Omega,\alpha}]f\right\|_{LM_{q_1,\varphi_2}^{\{x_0\}}} &= \sup_{r>0} \varphi_2(x_0,r)^{-1} |B(x_0,r)|^{-\frac{1}{q_1}} \left\|[\overrightarrow{b}, T_{\Omega,\alpha}]f\right\|_{L_{q_1}(B(x_0,r))} \\
&\leq C\prod_{i=1}^m \|\overrightarrow{b}\|_{LC_{p_i,\lambda_i}^{\{x_0\}}} \sup_{r>0} \varphi_2(x_0,r)^{-1} \\
&\quad \times \int_r^\infty \left(1+\ln\frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1}+\left(\sum_{i=1}^m \lambda_i+\sum_{i=1}^m \frac{1}{p_i}\right)\right)-1} \|f\|_{L_p(B(x_0,t))}dt \\
&\leq C\prod_{i=1}^m \|\overrightarrow{b}\|_{LC_{p_i,\lambda_i}^{\{x_0\}}} \|f\|_{LM_{p,\varphi_1}^{\{x_0\}}}.
\end{aligned}$$

For the case of $q_1 < s$, we can also use the same method, so we omit the details. Thus, we finish the proof of Theorem 1. □

Corollary 2. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $0 < \alpha < n$ and $1 < q, q_1, p_i, p < \frac{n}{\alpha}$ with $\frac{1}{q} = \sum_{i=1}^m \frac{1}{p_i} + \frac{1}{p}$, $\frac{1}{q_1} = \frac{1}{q} - \frac{\alpha}{n}$ and $\overrightarrow{b} \in LC_{p_i,\lambda_i}^{\{x_0\}}(\mathbb{R}^n)$ for $0 \leq \lambda_i < \frac{1}{n}$, $i = 1, \ldots, m$. Let also, for $s' \leq q$ the pair (φ_1, φ_2) satisfy condition (2.1) and for $q_1 < s$ the pair (φ_1, φ_2) satisfy condition (2.2). Then, the operators $M_{\Omega,\overrightarrow{b},\alpha}$ and $[\overrightarrow{b}, \overline{T}_{\Omega,\alpha}]$ are bounded from $LM_{p,\varphi_1}^{\{x_0\}}$ to $LM_{q_1,\varphi_2}^{\{x_0\}}$.*

Corollary 3. [13, 16] *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $T_{\Omega,\alpha}$ be a linear operator satisfying condition (1.7) and bounded from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$. Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $b \in LC^{\{x_0\}}_{p_2,\lambda}(\mathbb{R}^n)$, $0 \le \lambda < \frac{1}{n}$, $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, $\frac{1}{q_1} = \frac{1}{p_1} - \frac{\alpha}{n}$.*

Let also, for $s' \le p$ the pair (φ_1, φ_2) satisfy the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p_1}}}{t^{\frac{n}{q_1}+1-n\lambda}} dt \le C\, \varphi_2(x_0, r),$$

and for $q_1 < s$ the pair (φ_1, φ_2) satisfy the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname*{essinf}_{t<\tau<\infty} \varphi_1(x_0,\tau)\tau^{\frac{n}{p_1}}}{t^{\frac{n}{q_1}-\frac{n}{s}+1-n\lambda}} dt \le C\, \varphi_2(x_0, r) r^{\frac{n}{s}},$$

where C does not depend on r.

Then, the operator $[b, T_{\Omega,\alpha}]$ is bounded from $LM^{\{x_0\}}_{p_1,\varphi_1}$ to $LM^{\{x_0\}}_{q,\varphi_2}$. Moreover,

$$\left\|[b, T_{\Omega,\alpha}]\right\|_{LM^{\{x_0\}}_{q,\varphi_2}} \lesssim \|b\|_{LC^{\{x_0\}}_{p_2,\lambda}} \|f\|_{LM^{\{x_0\}}_{p_1,\varphi_1}}.$$

Theorem 2. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $T_{\Omega,\alpha}$ be a linear operator satisfying condition (1.7). Let also $0 < \alpha < n$ and $1 < q, q_1, p_i, p < \frac{n}{\alpha}$ with $\frac{1}{q} = \sum_{i=1}^m \frac{1}{p_i} + \frac{1}{p}$, $\frac{1}{q_1} = \frac{1}{q} - \frac{\alpha}{n}$ and $\overrightarrow{b} \in LC^{\{x_0\}}_{p_i,\lambda_i}(\mathbb{R}^n)$ for $0 \le \lambda_i < \frac{1}{n}$, $i = 1, \dots, m$. Let for $s' \le q$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6) and*

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m \varphi_1(x_0,t) \frac{t^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1} - \left(\sum_{i=1}^m \lambda_i + \sum_{i=1}^m \frac{1}{p_i}\right)\right)+1}} \le C_0\, \varphi_2(x_0, r), \tag{2.5}$$

where C_0 does not depend on $r > 0$,

$$\lim_{r\to 0} \frac{\ln \frac{1}{r}}{\varphi_2(x_0, r)} = 0 \tag{2.6}$$

and

$$c_\delta := \int_\delta^\infty (1 + \ln|t|)^m \varphi_1(x_0,t) \frac{t^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1} - \left(\sum_{i=1}^m \lambda_i + \sum_{i=1}^m \frac{1}{p_i}\right)\right)+1}} dt < \infty \tag{2.7}$$

for every $\delta > 0$, and for $q_1 < s$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6) and also

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m \varphi_1(x_0,t) \frac{t^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1} - \left(\frac{1}{s} + \sum_{i=1}^m \lambda_i + \sum_{i=1}^m \frac{1}{p_i}\right)\right)+1}} \le C\, \varphi_2(x_0, r) r^{\frac{n}{s}}, \tag{2.8}$$

where C_0 does not depend on $r>0$,

$$\lim_{r\to 0}\frac{\ln\frac{1}{r}}{\varphi_2(x_0,r)}=0$$

and

$$c_{\delta'}:=\int_{\delta'}^{\infty}\left(1+\ln|t|\right)^m\varphi_1(x_0,t)\frac{t^{\frac{n}{p}}}{t^{n\left(\frac{1}{q_1}-\left(\frac{1}{s}+\sum_{i=1}^m\lambda_i+\sum_{i=1}^m\frac{1}{p_i}\right)\right)+1}}dt<\infty \tag{2.9}$$

for every $\delta'>0$.

Then the operator $[\overrightarrow{b},T_{\Omega,\alpha}]$ is bounded from $VLM^{\{x_0\}}_{p,\varphi_1}$ to $VLM^{\{x_0\}}_{q_1,\varphi_2}$. Moreover,

$$\left\|[\overrightarrow{b},T_{\Omega,\alpha}]f\right\|_{VLM^{\{x_0\}}_{q_1,\varphi_2}}\lesssim\prod_{i=1}^m\|\overrightarrow{b}\|_{LC^{\{x_0\}}_{p_i,\lambda_i}}\|f\|_{VLM^{\{x_0\}}_{p,\varphi_1}}. \tag{2.10}$$

Proof. Since the inequality (2.10) holds by Theorem 1, we only have to prove the implication

(2.11)

$$\lim_{r\to 0}\frac{r^{-\frac{n}{p}}\|f\|_{L_p(B(x_0,r))}}{\varphi_1(x_0,r)}=0 \text{ implies } \lim_{r\to 0}\frac{r^{-\frac{n}{q_1}}\left\|[\overrightarrow{b},T_{\Omega,\alpha}]f\right\|_{L_{q_1}(B(x_0,r))}}{\varphi_2(x_0,r)}=0.$$

To show that

$$\frac{r^{-\frac{n}{q_1}}\left\|[\overrightarrow{b},T_{\Omega,\alpha}]f\right\|_{L_{q_1}(B(x_0,r))}}{\varphi_2(x_0,r)}<\epsilon \text{ for small } r,$$

we use the estimate (1.15):

$$\frac{r^{-\frac{n}{q_1}}\left\|[\overrightarrow{b},T_{\Omega,\alpha}]f\right\|_{L_{q_1}(B(x_0,r))}}{\varphi_2(x_0,r)}\lesssim\frac{\prod_{i=1}^m\|\overrightarrow{b}\|_{LC^{\{x_0\}}_{p_i,\lambda_i}}}{\varphi_2(x_0,r)}\int_r^{\infty}\left(1+\ln\frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1}+\left(\sum_{i=1}^m\lambda_i+\sum_{i=1}^m\frac{1}{p_i}\right)\right)-1}\|f\|_{L_p(B(x_0,t))}dt.$$

We take $r<\delta_0$, where δ_0 is small enough and split the integration:

$$\frac{r^{-\frac{n}{q_1}}\left\|[\overrightarrow{b},T_{\Omega,\alpha}]f\right\|_{L_{q_1}(B(x_0,r))}}{\varphi_2(x_0,r)}\le C\left[I_{\delta_0}(x_0,r)+J_{\delta_0}(x_0,r)\right], \tag{2.12}$$

where $\delta_0>0$ (we may take $\delta_0<1$), and

$$I_{\delta_0}(x_0,r):=\frac{1}{\varphi_2(x_0,r)}\int_r^{\delta_0}\left(1+\ln\frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1}+\left(\sum_{i=1}^m\lambda_i+\sum_{i=1}^m\frac{1}{p_i}\right)\right)-1}\|f\|_{L_p(B(x_0,t))}dt,$$

and

$$J_{\delta_0}(x_0,r):=\frac{1}{\varphi_2(x_0,r)}\int_{\delta_0}^{\infty}\left(1+\ln\frac{t}{r}\right)^m t^{n\left(-\frac{1}{q_1}+\left(\sum_{i=1}^m\lambda_i+\sum_{i=1}^m\frac{1}{p_i}\right)\right)-1}\|f\|_{L_p(B(x_0,t))}dt$$

and $r < \delta_0$. Now we can choose any fixed $\delta_0 > 0$ such that

$$\frac{t^{-\frac{n}{p}} \|f\|_{L_p(B(x_0,t))}}{\varphi_1(x_0,t)} < \frac{\epsilon}{2CC_0}, \qquad t \leq \delta_0,$$

where C and C_0 are constants from (2.5) and (2.12). This allows to estimate the first term uniformly in $r \in (0, \delta_0)$:

$$CI_{\delta_0}(x_0,r) < \frac{\epsilon}{2}, \qquad 0 < r < \delta_0.$$

For the second term, writing $1 + \ln \frac{t}{r} \leq 1 + |\ln t| + \ln \frac{1}{r}$, we obtain

$$J_{\delta_0}(x_0,r) \leq \frac{c_{\delta_0} + \widetilde{c_{\delta_0}} \ln \frac{1}{r}}{\varphi_2(x_0,r)} \|f\|_{LM^{\{x_0\}}_{p,\varphi_1}},$$

where c_{δ_0} is the constant from (2.7) with $\delta = \delta_0$ and $\widetilde{c_{\delta_0}}$ is a similar constant with omitted logarithmic factor in the integrand. Then, by (2.6) we can choose small enough r such that

$$J_{\delta_0}(x_0,r) < \frac{\epsilon}{2},$$

which completes the proof of (2.11).

For the case of $q_1 < s$, we can also use the same method, to obtain the desired result. Therefore, the proof of Theorem 2 is completed. □

Remark 2. *Conditions (2.7) and (2.9) are not needed in the case when $\varphi(x_0, r)$ does not depend on x_0, since (2.7) follows from (2.5) and similarly, (2.9) follows from (2.8) in this case.*

Corollary 4. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $0 < \alpha < n$ and $1 < q, q_1, p_i, p < \frac{n}{\alpha}$ with $\frac{1}{q} = \sum_{i=1}^{m} \frac{1}{p_i} + \frac{1}{p}$, $\frac{1}{q_1} = \frac{1}{q} - \frac{\alpha}{n}$ and $\overrightarrow{b} \in LC^{\{x_0\}}_{p_i,\lambda_i}(\mathbb{R}^n)$ for $0 \leq \lambda_i < \frac{1}{n}$, $i = 1, \ldots, m$. Let also, for $s' \leq q$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6)-(2.6) and (2.5)-(2.7) and for $q_1 < s$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6)-(2.6) and (2.8)-(2.9). Then, the operators $M_{\Omega,\overrightarrow{b},\alpha}$ and $[\overrightarrow{b}, \overline{T}_{\Omega,\alpha}]$ are bounded from $VLM^{\{x_0\}}_{p,\varphi_1}$ to $VLM^{\{x_0\}}_{q_1,\varphi_2}$.*

Corollary 5. *Suppose that $x_0 \in \mathbb{R}^n$, $\Omega \in L_\infty(\mathbb{R}^n) \times L_s(S^{n-1})$, $s > 1$, is homogeneous of degree zero with respect to the second variable y on $\mathbb{R}^n$. Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $b \in LC^{\{x_0\}}_{p_2,\lambda}(\mathbb{R}^n)$, $0 \leq \lambda < \frac{1}{n}$, $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, $\frac{1}{q_1} = \frac{1}{p_1} - \frac{\alpha}{n}$, and $T_{\Omega,\alpha}$ is a linear operator satisfying condition (1.7) and bounded from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$. Let also, for $s' \leq p$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6) and*

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x_0,t) \frac{t^{\frac{n}{p_1}}}{t^{\frac{n}{q_1}+1-n\lambda}} dt \leq C_0 \varphi_2(x_0,r),$$

where C_0 does not depend on $r > 0$,

$$\lim_{r\to 0} \frac{\ln \frac{1}{r}}{\varphi_2(x_0,r)} = 0$$

and

$$c_\delta := \int_\delta^\infty (1 + \ln|t|) \varphi_1(x_0,t) \frac{t^{\frac{n}{p_1}}}{t^{\frac{n}{q_1}+1-n\lambda}} dt < \infty$$

for every $\delta > 0$, and for $q_1 < s$ the pair (φ_1, φ_2) satisfies conditions (1.5)-(1.6) and also

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x_0, t) \frac{t^{\frac{n}{p_1}}}{t^{\frac{n}{q_1} - \frac{n}{s} + 1 - n\lambda}} dt \leq C_0 \varphi_2(x_0, r) r^{\frac{n}{s}},$$

where C_0 does not depend on $r > 0$,

$$\lim_{r \to 0} \frac{\ln \frac{1}{r}}{\varphi_2(x_0, r)} = 0$$

and

$$c_{\delta'} := \int_{\delta'}^\infty (1 + \ln |t|) \varphi_1(x_0, t) \frac{t^{\frac{n}{p_1}}}{t^{\frac{n}{q_1} - \frac{n}{s} + 1 - n\lambda}} dt < \infty$$

for every $\delta' > 0$.

Then the operator $[b, T_{\Omega,\alpha}]$ is bounded from $VLM^{\{x_0\}}_{p_1,\varphi_1}$ to $VLM^{\{x_0\}}_{q,\varphi_2}$. Furthermore, we have

$$\|[b, T_{\Omega,\alpha}] f\|_{VLM^{\{x_0\}}_{q,\varphi_2}} \lesssim \|b\|_{LC^{\{x_0\}}_{p_2,\lambda}} \|f\|_{VLM^{\{x_0\}}_{p_1,\varphi_1}}.$$

References

[1] A. Abdalmonem, O. Abdalrhman, S. Tao, The boundedness of fractional integral with variable kernel on variable exponent Herz-Morrey spaces, J. Appl. Math. Phys., 4 (2016), 787-795.

[2] A. Abdalmonem, O. Abdalrhman, S. Tao, Boundedness of fractional integral with variable kernel and their commutators on variable exponent Herz spaces, Appl. Math. 7 (2016), 1165-1182.

[3] A.S. Balakishiyev, V.S. Guliyev, F. Gurbuz and A. Serbetci, Sublinear operators with rough kernel generated by Calderon-Zygmund operators and their commutators on generalized local Morrey spaces, J. Inequal. Appl. 2015, 2015:61. doi:10.1186/s13660-015-0582-y.

[4] L. Caffarelli, Elliptic second order equations, Rend. Semin. Math. Fis. Milano, 58 (1990), 253-284.

[5] X.N. Cao, D.X. Chen, The boundedness of Toeplitz-type operators on vanishing Morrey spaces. Anal. Theory Appl. 27 (2011), 309-319.

[6] S. Chanillo, A note on commutators, Indiana Univ. Math. J., 31 (1), (1982), 7-16.

[7] Y. Chen, Y. Ding, R. Li, The boundedness for commutator of fractional integral operator with rough variable kernel, Potential Anal., 38 (2013), 119-142.

[8] F. Chiarenza, M. Frasca, P. Longo, Interior $W^{2,p}$-estimates for nondivergence elliptic equations with discontinuous coefficients, Ricerche Mat., 40 (1) (1991), 149-168.

[9] F. Chiarenza, M. Frasca, P. Longo, $W^{2,p}$-solvability of Dirichlet problem for nondivergence elliptic equations with VMO coefficients, Trans. Amer. Math. Soc., 336 (2) (1993), 841-853.

[10] R.R. Coifman, R. Rochberg, G. Weiss, Factorization theorems for Hardy spaces in several variables, Ann. Math., 103 (3) (1976), 611-635.

[11] R.R. Coifman, P.L. Lions, Y. Meyer, S. Semmes, Compensated compactness and Hardy spaces, J. Math. Pure Appl., 72 (3) (1993), 247-286.

[12] G. Di Fazio, D.K. Palagachev and M.A. Ragusa, Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients, J. Funct. Anal., 166 (1999), 179-196.

[13] F. Gurbuz, Boundedness of some potential type sublinear operators and their commutators with rough kernels on generalized local Morrey spaces [*Ph.D. thesis*], Ankara University, Ankara, Turkey, 2015 (in Turkish).

[14] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic Calderón-Zygmund operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, Open Math., 14 (2016), 300-323.

[15] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic fractional integral operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, Adv. Math. (China), 2016, in press.

[16] F. Gurbuz, Sublinear operators with a rough kernel generated by fractional integrals and local Campanato space estimates for commutators with rough kernel on generalized local Morrey spaces, Int. J. Appl. Math. & Stat., 2016, in press.
[17] S. Janson, Mean oscillation and commutators of singular integral operators, Ark. Mat., 16 (1978), 263-270.
[18] G. Lu, S.Z. Lu, D.C. Yang, Singular integrals and commutators on homogeneous groups, Anal. Math., 28 (2002), 103-134.
[19] S.Z. Lu and D.C. Yang, The central BMO spaces and Littlewood-Paley operators, Approx. Theory Appl. (N.S.), 11 (1995), 72-94.
[20] C. Miranda, Sulle equazioni ellittiche del secondo ordine di tipo non variazionale, a coefficienti discontinui. Ann. Math. Pura E Appl. 63 (4) (1963), 353-386.
[21] A. Mazzucato, Besov-Morrey spaces:functions space theory and applications to non-linear PDE., Trans. Amer. Math. Soc., 355 (2002), 1297-1364.
[22] H. Mo, H.Y. Xue, Commutators generated by singular integral operators with variable kernels and local Campanato functions on generalized local Morrey spaces, Commun. Math. Anal., 19 (2) (2016), 32-42.
[23] C.B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, Trans. Amer. Math. Soc., 43 (1938), 126-166.
[24] B. Muckenhoupt, R.L. Wheeden, Weighted norm inequalities for singular and fractional integrals. Trans. Amer. Math. Soc., 161 (1971), 249-258.
[25] D.K. Palagachev, L.G. Softova, Singular integral operators, Morrey spaces and fine regularity of solutions to PDE's, Potential Anal., 20 (2004), 237-263.
[26] M. Paluszynski, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss, Indiana Univ. Math. J., 44 (1995), 1-17.
[27] L.E. Persson, M.A. Ragusa, N. Samko, P. Wall, Commutators of Hardy operators in vanishing Morrey spaces. AIP Conf. Proc. 1493, 859 (2012); http://dx.doi.org/10.1063/1.4765588.
[28] M.A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces. J. Global Optim. 368 (40) (2008), 1-3.
[29] A. Ruiz, L. Vega, On local regularity of Schrödinger equations, Int. Math. Res. Not., 1 (1993), 13-27.
[30] S.G. Shi, S.Z. Lu, A characterization of Campanato space via commutator of fractional integral, J. Math. Anal. Appl., 419 (2014), 123-137.
[31] L.G. Softova, Singular integrals and commutators in generalized Morrey spaces, Acta Math. Sin. (Engl. Ser.), 22 (2006), 757-766.
[32] F. Soria, G. Weiss, A remark on singular integrals and power weights, Indiana Univ. Math. J., 43 (1994) 187-204.
[33] C. Vitanza, Functions with vanishing Morrey norm and elliptic partial differential equations. In: Proceedings of Methods of Real Analysis and Partial Differential Equations,Capri, pp. 147-150. Springer (1990).
[34] C. Vitanza, Regularity results for a class of elliptic equations with coefficients in Morrey spaces. Ricerche Mat., 42 (2) (1993), 265-281.
[35] H. Wang, Singular and fractional integral operators with variable kernels on the weak Hardy spaces, Commun. Math. Anal., 18 (1) (2015), 48-63.
[36] R.L. Wheeden and A. Zygmund, Measure and Integral: An Introduction to Real Analysis, vol. 43 of Pure and Applied Mathematics, Marcel Dekker, New York, NY, USA, 1977.
[37] P. Zhang and K. Zhao, Commutators of integral operators with variable kernels on Hardy spaces, Proc. Indian Acad. Sci. (Math. Sci.), 115 (4) (2003), 399-410.

ANKARA UNIVERSITY, FACULTY OF SCIENCE, DEPARTMENT OF MATHEMATICS, TANDOĞAN 06100, ANKARA, TURKEY
E-mail address: feritgurbuz84@hotmail.com